



ACF

Arctic Climate Forum

November 2019 – April 2020

Arctic Seasonal Review

Vasily Smolyanitsky

Anna Danshina, Anastassiya Revina

Arctic and Antarctic Research Institute, Russia

with contributions from: Copernicus CCS, AARI (Review of hydrometeorological processes in Northern Polar Region in 2019), DMI PolarPortal, SnowWatch (FMI, ECCC, Rutgers Glob Snow Lab)



WMO OMM

World Meteorological Organization

Organisation météorologique mondiale

Content of review

- ❖ Review for 2 periods: NDJ 2019/2020 and FMA 2020
- ❖ Atmosphere variables include:
 - ✓ Atmospheric circulation (MSL) and geopotential height (gp50, gp500)
 - ✓ Surface air temperature
 - ✓ Precipitation
- ❖ Sea ice variables include:
 - ✓ atmosphere and polar ocean precursors
 - ✓ Ice extent and ice conditions analysis
 - ✓ Sea ice thickness and volume reanalysis
- ❖ Polar Ocean
 - ✓ SST, waves and swell height (storminess)
 - ✓ pH (acidification/alkalization estimates)
- ❖ Solid precipitation (land snow)
- ❖ Briefs on current status (SAT, winds, Prec, sea ice, snow)




Atmosphere:


- ✓ Precursors - atmospheric circulation patterns
- ✓ Surface air temperature
- ✓ Precipitation



NDJ 2019/2020 atmospheric circulation



MSLP hPa anomalies (norms 1981-2010)




H50 ranks (1979-2020)


- ❖ Based on atmosphere numerical model (reanalysis)
- ❖ Negative mean sea level atmospheric pressure (MSLP) anomalies (lower pressure, marked in blue) dominated through the European, Siberian and Canadian archipelago regions
- ❖ Opposite situation (higher pressure, marked in red) was observed over Svalbard, Canadian and Alaska regions, Bering Sea
- ❖ That led to prevalence of zonal form of circulation (transfer of heat/cold west/east) in the troposphere with the center of polar vortex over western Siberia as seen on the 50 hPa geopotential height (H50)

AARI / ERA5 reanalysis

FMA 2020 atmospheric circulation



MSLP hPa anomalies (norms 1981-2010)




H50 (left) and H500 ranks (1979-2020)

- ❖ Much stronger negative MSLP anomalies (lower pressure, marked in blue) were observed during FMA 2020 in the European region, Western Siberia, Arctic Ocean but not in Alaska and Bering Sea
- ❖ That again led to increased cyclonic activity, more polar lows with further increased precipitation
- ❖ Polar vortex was very intense as observed at H50 pattern and caused meridian type of circulation with several 'heat waves' in Western and Eastern Siberia

[AARI / ERA5 reanalysis]



December 2019 – February 2020 SAT (T2m): anomalies and ranks (observation)



- ❖ The winter air temperature across Arctic was above normal except Alaska, Greenland, Svalbard, some parts of Canadian archipelago and Chukchi region.
- ❖ The most notable positive anomalies were present across of Western and Eastern Siberia Alaska and some parts of N Atlantic
- ❖ Very close to record high temperatures were observed in Eastern Siberia.









[AARI]



SAT anomalies by regions during winter 2019/20 (DJF) (observations)

Region	Anomaly	Rank	The warmest year (anomaly)	The coldest year (anomaly)
North Atlantic	1,9	14	2014 (3,9)	1966 (-2,4)
Barents	5,4	3	1937 (6,5)	1979 (-4,4)
Western Siberia	7,2	3	2012, 2016 (7,6)	1969 (-5,6)
Eastern Siberia	4,6	2	2016 (4,6)	1966 (-4,5)
Chukchi	1,8	13	2018 (6,7)	2002 (-2,3)
Western Canada	-0,9	42	2018 (6,1)	1965 (-5,6)
Eastern Canada	1,2	18	2010 (5,0)	1972 (-3,6)

Reference period: 1961-1990











[AARI]

SAT anomalies by Arctic seas during winter 2019/20 (DJF) (observations)

Sea	Anomaly	Rank	The warmest year (anomaly)	The coldest year (anomaly)
Northern part of Greenland and Norwegian Seas	2,1	19	2014 (6,1)	1966 (-2,7)
Barents Sea	3,3	20	1937 (7,6)	1979 (-3,4)
Kara Sea	6,5	8	1945 (9,8)	1979 (-5,9)
Laptev Sea	4,4	3	2012 (4,6)	1979 (-4,4)
Eastern Siberian Sea	2,9	7	2018 (6,5)	1966 (-3,3)
Chukchi Sea	1,6	19	2018 (8,8)	1939 (-3,4)
Beaufort Sea	0,7	29	2018 (6,3)	1966 (-3,1)
N part of Canadian arhipelago	1,0	18	2010 (5,0)	1949 (-3,6)


Reference period: 1961-1990




[AARI]

NDJ 2019-2020, FMA 2020 – SAT (T2m): anomalies and ranks

NDJ 2019/2020




SAT anomaly, 1981-2010




SAT rank, 1979-2020

FMA 2020



SAT anomaly, 1981-2010



Rank, 1979-2020









❖ For the whole season from November 2019 – April 2020 positive close to maximum air temperature anomalies prevailed over Western and Eastern Siberia, with negative anomalies prevailing in marine Barents, Alaska and parts of Western Canada

❖ Atlantic and partly Chukchi areas experienced switch from positive to negative anomalies during NDJ - FMA

AARI / ERA5 reanalysis

Precipitation trends for ND 2019 - J 2020

- ❖ Analysis based on observations by the Arctic seas
- ❖ General positive trends – wetter conditions – for the Nordic seas, Beaufort Sea
- ❖ General negative trends – drier conditions - for Siberian shelf seas
- ❖ No general significant trends for Canadian Arctic regions




Reference period: 1961-1990

[AARI]




Precipitation (Prec) NDJ and FMA 2019/2020: anomalies and ranks

NDJ 2019/2020




Prec Anomaly, 1981-2010




Prec Rank, 1979-2020

FMA 2020



Prec Anomaly, 1981-2010



Prec Rank, 1979-2020

- ❖ For NDJ 2019/2020 and FMA 2020 Barents and Western Siberia regions saw very wet seasons
- ❖ Same wetter conditions observed for Alaska during FMA 2020
- ❖ Drier conditions were observed over Svalbard, parts of Greenland, Sea of Okhotsk

[AARI / ERA5 reanalysis]




Sea ice variables:


- ✓ Precursors in atmosphere and polar ocean
- ✓ Ice extent and ice conditions analysis
- ✓ Sea ice thickness and volume based on coastal stations and reanalysis




Atmosphere – polar ocean precursors for winter – spring 2019 – 2020 sea ice conditions




SAT SON anomaly,
1981-2010



SAT DJF anomaly,
1981-2010







Heat Capacity anomaly OND,
2000-2019



Heat Capacity anomaly JFM,
2000-2019


- ❖ High positive anomalies in surface air temperature (SAT) as well as prevailing positive polar ocean heat capacity (HC) in upper 15m during OND 2019 slowed in general freeze-up and sea ice growth in the Arctic
- ❖ Further in time lesser positive SAT anomalies as well as in general neutral HC anomalies during JFM stimulated ice extent growth
- ❖ Prominent negative HC anomalies lead to close to normal ice growth in the N Barents Sea and Sea of Okhotsk

Arctic (NH) seasonal ice extent 1978.... 2020



S, 1000 km ² *
2017 14467
2018 14516
2015 14526
2016 14580
2011 14701
2006 14867
2019 14891
2007 14931
2014 14972
2005 15101
2020 15159
...
1988 16461
1983 16547
1979 16769


* - maximum for Mar (AARI)



❖ Maximum winter ice extent, 11th in row, 15,16 mln km² (14,89 in 2019) reached 4 March 2020 (11 March in 2019)

❖ During freezing period (ND 2019) lowest on record was observed due to extreme minimums in Bering, Chukchi Seas


Seasonal NH ice extent variability: 1978 - 2020



- ❖ Seasonal patterns of daily ice extent allows to analysis seasonal variability of ice extent
- ❖ Both winter maximums and summer minimums continue to diminish
- ❖ Though, significant interannual variability of ice extent occurs




NDJ 2019/2020 Arctic sea ice – concentration and stage of development



18-21 Nov

16-19 Dec


13-16 Jan




Blended AARI/CIS/NIC (JCOMM) ice charts; ice edge – nearest 5days, reference period: 1999-2018

Weather • Climate • Water


FMA 2020 Arctic sea ice – concentration and stage of development






17-21 Feb



16-19 Mar




13-17 Apr



Blended AARI/CIS/NIC (JCOMM) ice charts; ice edge – nearest 5 days, reference period: 1999-2018

Sea ice fast ice maximum thickness values and anomalies by end of April/Mar 2020 (stations)



WMO stations used:

Russia: 12 (Varandey, Amderma, Belyi, Dikson, Sterlegova, Cheluskin, Tiksi, Kotelnnyi, Sannikova, Ayon, Valkarkay)



- ❖ Observed maximum winter ice thicknesses significantly less than normal (for the last 30 years) for Kara Sea (up to -50 cm, which is opposite to 2019) and slightly less than normal for Eastern Siberian Sea
- ❖ Slightly thicker ice observed in Laptev Sea

Ref [1991-2020]



[AARI]

Sea ice thickness for 15 Mar 2004...2020 and ice volume




- ❖ Models show Arctic ice volume in 2020 the lowest from Oct 2019 for 2004-2020 with 2nd in row in 2016
- ❖ The present ice extent slightly higher then in 2016 means sea ice thicknesses in winter / spring 2019/2020 are in general lower then in 2019

Polar Ocean:


- ✓ Sea surface temperature
- ✓ pH and acidification or alkalization of the Arctic ?
- ✓ Storms - Wave and swell height




Waves and ph in the Arctic Ocean - NDJ 2019-2020, FMA 2020




SST NDJ anomaly, 1981-2010




SST FMA anomaly, 1981-2010




WW&S height NDJ rank, 1979-2020



WW&S height FMA rank, 1979-2020



pH anomaly 2m SON 2000-2019



pH anomaly 2m DJF 2000-2019

- ❖ Boundary seas of the Arctic Ocean were in general warmer and stormy during winter-spring 2019-2020 with exceptions Svalbard and N Greenland (colder, calmer), Sea of Okhotsk (colder)
- ❖ Numerical models show both positive (Arctic Basin, Chukchi Sea) and negative pH (Barents, Kara Sea, Canadian Arctic) anomalies to the last 20 years, that allows occurrence of both alkalization and acidification processes in the Arctic (no effect to wildlife?)

AARI / Copernicus Climate Change Service
(ERA5 & MERCATOR reanalysis)

Land Snow:


- ✓ Snow water equivalent
- ✓ Snow extent




Land snow (satellite, obs)

- ❖ Snow extent in winter-spring 2019/2020 was less than normal with prominent negative anomalies (no snow) in most of European sector
- ❖ Positive anomalies (more snow) were observed in Scandinavia, southern Canada
- ❖ Greater Snow Water Equivalent in 2019/2020 means higher snow height observed


Departure from Normal - November 2019




Departure from Normal - December 2019




Departure from Normal - January 2020




Departure from Normal - February 2020






Departure from Normal - March 2020



Departure from Normal - April 2020




Ref [1981-2010]




[FMI, ECCC, Rutgers Glob Snow Lab / GCW]]

Current Conditions (21-26 May 2020)

May 23 to May 27, 2020




May 23 to May 27, 2020




SAT, precipitation, mean wind vectors, NAO for 23-27.05.2020 (<http://polarportal.dk>)

- ❖ Till now week westerly, moderate northern winds in European sector and strong southern winds over Siberia (NAO~0) led to lower SAT in European sector, prominent higher SAT over Siberia and the Arctic Ocean ('heat waves')
- ❖ Northern Scandinavia, Arctic coasts, Chukchi peninsula are still under snow
- ❖ NE part of Barents Sea, Kara, W Laptev Seas, Chukchi Sea are under intense ice melt which is extreme
- ❖ However general pattern of TransArctic drift keeps ice conditions in Greenland Sea and Svalbard waters near normal



Daily Snow - May 27, 2020 (Day 148)



Snow extent for 27 May 2020, Rutgers Global snow lab

**Thank you! Merci! Takk! Спасибо!
Tak! Tack! Kiitos! þakka þér fyrir!
Naqurmiik ! Qaġaasakuq !
Grazie! Giitu! Vielen Dank!
Dhanyavaad !**



WMO OMM

World Meteorological Organization

Organisation météorologique mondiale